



CERTIFICATE OF ANALYSIS

PRODUCT NAME: GLYKO® $\beta(1-4,6)$ GALACTOSIDASE (Jack Bean)
PRODUCT CODE: GKX-5012
LOT NUMBER: DG33 020a
FORMULATION: Lyophilized from 20 mM sodium citrate phosphate (pH 6.0)
RECONSTITUTION: Dissolve the lyophilizate in 169 μ l of high purity water to obtain the described formulation. Since the enzyme will be more concentrated than is required for most applications, dilute further with buffer as needed.
SUGGESTIONS FOR USE: Conditions for use vary depending on the application, since the enzyme hydrolyzes $\beta(1-4)$ and $\beta(1-6)$ linkages significantly more rapidly than $\beta(1-3)$ linkages. For example, for non-selective hydrolysis of non-reducing terminal galactose, a final concentration of at least 4 U/ml is recommended whereas a final concentration of <1 U/ml is recommended for selective hydrolysis of $\beta(1-4)$ - and $\beta(1-6)$ -linked galactose.
STORAGE: -20°C until redissolved. Store redissolved enzyme at $2-8^{\circ}\text{C}$ or -20°C but avoid repeated freeze-thaw cycles.
PACK SIZE: 5 Units
EXPIRATION: August 2018

QUALITY CONTROL

1.	Specific activity ¹ :	Passed	(Specification: ≥ 70 U/mg)
2.	Protease assay ² :	Passed	(Specification: "Not Detectable")
3.	Contaminants ³ : (except as noted below)	Passed	(Specification: $\leq 0.001\%$)
	α -Galactosidase	0.0061%	
	β -Glucosidase	0.0462%	
	β -Xylosidase	0.0039%	

Authorized Signature

1. One unit of Jack Bean β -Galactosidase is defined as the amount of enzyme required to catalyze the release of one μ mole of p-nitrophenol per minute from pNP- β -galactopyranoside at pH 3.5 and 37°C .
2. No protease activity was detectable after incubation of the enzyme with 0.2 mg resorufin-labeled casein for ~18 hours at 37°C based on Schickaneder E, Hösel W, von der Eltz H, Geuß U. Casein-resorufin, a new substrate for a highly sensitive protease assay. Fresenius Z. Anal Chem. 1988 330:360.
3. The absence of exoglycosidase contaminants was confirmed by extended incubations with the corresponding pNP-glycosides: α -fucosidase, α -mannosidase, β -mannosidase, β -N-acetylhexosaminidase, α -N-acetylglactosaminidase, α -galactosidase, α -glucosidase, β -glucosidase and β -xylosidase. The absence of contaminating sialidase was confirmed by extended incubation with MU-NANA. **Note: this enzyme has activity on pNP- β -D-fucoside (although reduced relative to the standard substrate).**