



CERTIFICATE OF ANALYSIS

PRODUCT NAME: GLYKO® β (1-4,6)GALACTOSIDASE (Jack Bean)

PRODUCT CODE: GKX-5012

LOT NUMBER: DG33 019a

FORMULATION: Lyophilized from 20 mM sodium citrate phosphate (pH 6.0)

RECONSTITUTION: Dissolve the lyophilizate in 124 μ l of high purity water to obtain the described formulation. Since the enzyme will be more concentrated than is required for most applications, dilute further with buffer as needed.

SUGGESTIONS FOR USE: Conditions for use vary depending on the application, since the enzyme hydrolyzes β (1-4) and β (1-6) linkages significantly more rapidly than β (1-3) linkages. For example, for non-selective hydrolysis of non-reducing terminal galactose, a final concentration of at least 4 U/ml is recommended whereas a final concentration of <1 U/ml is recommended for selective hydrolysis of β (1-4)- and β (1-6)-linked galactose.

STORAGE: -20°C until redissolved. Store redissolved enzyme at 2-8°C or -20°C but avoid repeated freeze-thaw cycles.

PACK SIZE: 5 Units

EXPIRATION: July 2018 (extended from prior exp. date based on re-assay)

RE-ASSAY DATE: October 2017

QUALITY CONTROL

1.	Specific activity ¹ :	Passed	(Specification: ≥ 70 U/mg)
2.	Protease assay ² :	Passed	(Specification: "Not Detectable")
3.	Contaminants ³ : (except as noted below)	Passed	(Specification: $\leq 0.001\%$)
	α -Galactosidase	0.007%	
	β -Glucosidase	0.037%	
	β -Xylosidase	0.003%	

Authorized Signature

1. One unit of Jack Bean β -Galactosidase is defined as the amount of enzyme required to catalyze the release of one μ mole of p-nitrophenol per minute from pNP- β -galactopyranoside at pH 3.5 and 37°C.
2. No protease activity was detectable after incubation of the enzyme with 0.2 mg resorufin-labeled casein for ~18 hours at 37°C based on Schickaneder E, Hösel W, von der Eltz H, Geuß U. Casein-resorufin, a new substrate for a highly sensitive protease assay. Fresenius Z. Anal Chem. 1988 330:360.

3. The absence of exoglycosidase contaminants was confirmed by extended incubations with the corresponding pNP-glycosides: α -fucosidase, α -mannosidase, β -mannosidase, β -N-acetylhexosaminidase, α -N-acetylgalactosaminidase, α -galactosidase, α -glucosidase, β -glucosidase and β -xylosidase. The absence of contaminating sialidase was confirmed by extended incubation with MU-NANA. **Note: this enzyme has activity on pNP- β -D-fucoside (although reduced relative to the standard substrate).**